

A solution of the Navier–Stokes system of equations in three-dimensions

R. K. Michael Thambynayagam†

Retired scientist, Schlumberger, Houston, Texas, USA

In this paper we derive a solution of the Navier-Stokes system of equations for incompressible fluid flows. We show that for a given smooth solenoidal initial velocity vector field there exist a spatially periodic solution of velocity and pressure that span the entire unbounded domain: $\mathbb{R}^3 = \{-\infty < x_i < \infty; i = 1, 2, 3\}$.

Key words: Navier-Stokes, Incompressible flow, Viscous flows, Euler flow, Partial differential equations

ORCID: <https://orcid.org/0000-0002-1778-7327>

1. Introduction

The Navier-Stokes equations are a system of partial differential equations that describe the flow of incompressible fluids: Navier (1827); Stokes (1845). These equations are non-linear in nature and therefore, difficult to solve analytically. Necessary background knowledge of the subject can be found in Batchelor (1967), Drazin and Riley (2007) and the monographs by Ladyzhenskaya (1969).

Ethier and Steinman (1994) presented unsteady analytical solutions involving all three Cartesian velocity components to the Navier-Stokes equations. Recently, Antuono (2020) extended Ethier and Steinman’s work to a class of solutions for the velocity vector field in a three-dimensional torus in terms of a wave number divided into two parts, characterized by positive and negative helicity respectively. In Appendix A, we provide Antuono’s solution with positive helicity.

In a previous article Thambynayagam (2013) presented a method to derive analytical solutions by recasting the Navier-Stokes equation in terms of three distinct terms associated, respectively, with the linear viscous forces, inertial forces and the external forces applied to the fluid. In this paper we apply the method to derive a single time evolution analytical solution for the velocity vector field and pressure that span the entire

† April 28, 2022. email: michael.thambynayagam@gmail.com

unbounded domain: $\mathbb{R}^3 = \{-\infty < x_i < \infty; i = 1, 2, 3\}$. Mathematical expressions, particularly equations (4.5)-(4.17), are tedious and exhaustively boring to derive, but they are straightforward.

2. The fundamental problem

In a Cartesian frame of reference, the fundamental equations of momentum and mass conservation for incompressible viscous fluid flow fields are represented as follows:

$$\frac{\partial v_i}{\partial t} + g_i = \kappa \Delta v_i - \frac{1}{\rho} \frac{\partial p}{\partial x_i} + f_i \quad x \in \mathbb{R}^n, \quad t \geq 0 \quad (2.1)$$

where $v(x, t) = v_i(x, t)$, $1 \leq i \leq n$, $x \in \mathbb{R}^n$ is the solenoidal velocity vector field and $p(x, t)$ the pressure of the fluid.

$$g_i = \sum_{j=1}^n v_j \frac{\partial v_i}{\partial x_j} \quad (2.2)$$

is the nonlinear inertial force, $f_i(x, t)$ are the components of an externally applied force, ρ is the constant density of the fluid, κ is the positive coefficient of kinematical viscosity and $\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$ is the Laplacian in the space variables.

The conservation of mass yields the divergence-free (incompressibility) condition

$$\mathbf{div} v = \sum_{i=1}^n \frac{\partial v_i}{\partial x_i} = 0 \quad x \in \mathbb{R}^n, \quad t \geq 0 \quad (2.3)$$

Conservation law also implies that the energy dissipation of a viscous fluid is bounded by the initial kinetic energy. Therefore, if the externally applied force does no net work on the fluid, the kinetic energy of the solution should be finite. There exist a constant E such that

$$\int_{\mathbb{R}^n} |\mathbf{v}|^2 dx < E \quad t \geq 0 \quad (2.4)$$

The Navier-Stokes equation (2.1) must be solved forward in time, starting from an initial divergence-free velocity field

$$v(x, 0) = v^{(0)}(x) \quad x \in \mathbb{R}^n \quad (2.5)$$

with the pressure evolving in time to maintain the incompressibility constraint (2.3). The superscript (0) is used to denote the value of a function at time zero.

3. Recasting the Navier-Stokes equation:

$$\mathbb{R}^n = \{-\infty < x_i < \infty; i = 1, 2, \dots, n\}$$

In this section, in order to be perspicuous, we repeat the material from Thambynayagam (2013) relating to the recasting of the Navier-Stokes equation and the resulting *lemma*. Assuming that the divergence and the linear operator can be commuted, the pressure field can be formally obtained by taking the divergence of (2.1) as a solution of the

Poisson equation, which is

$$\Delta p = \rho \sum_{i=1}^n \frac{\partial (f_i - g_i)}{\partial x_i} \quad (3.1)$$

Equation (3.1) is called the pressure Poisson equation. The use of pressure Poisson equation in solving the Navier-Stokes equation is discussed in a paper by Gresho and Sani (1987). It is important to note that while (2.1) and (2.3) lead to the pressure Poisson equation (3.1), the reverse; that is, (2.1) and (3.1), do not always lead to (2.3). In deriving solutions of the Navier-Stokes equations, we therefore ensure that the velocity vector field remains solenoidal at all times.

The general solution of the Poisson equation (3.1) is

$$\begin{aligned} p(x, t) &= -\frac{\rho}{2\pi} \int_{\mathbb{R}^2} P(y, t) \ln \left(\frac{1}{\sqrt{\mathcal{P}_n(x, y)}} \right) \prod_{j=1}^2 dy_j, \quad n = 2, \\ p(x, t) &= -\frac{\rho \Gamma(\frac{n}{2})}{2(n-2)\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n} \frac{P(y, t)}{\{\mathcal{P}_n(x, y)\}^{\frac{n-2}{2}}} \prod_{j=1}^n dy_j, \quad n \geq 3 \end{aligned} \quad (3.2)$$

where $\Gamma(z) = \int_0^\infty e^{-u} u^{z-1} du$ [$\Re z > 0$], is the Gamma function,

$$P(x, t) = \sum_{j=1}^n \frac{\partial (f_j - g_j)}{\partial x_j} \quad (3.3)$$

and

$$\mathcal{P}_n(x, y) = \sum_{j=1}^n (x_j - y_j)^2 \quad (3.4)$$

Differentiating (3.2) with respect to x_i we obtain

$$\frac{\partial p}{\partial x_i} = \frac{\rho \Gamma(\frac{n}{2})}{2\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n} \frac{(x_i - y_i) P(y, t)}{\{\mathcal{P}_n(x, y)\}^{\frac{n}{2}}} \prod_{j=1}^n dy_j, \quad n \geq 2 \quad (3.5)$$

Substituting for $\frac{\partial p}{\partial x_i}$ in (2.1) the following can be formulated:

$$\frac{\partial v_i}{\partial t} = \kappa \Delta v_i - \frac{\Gamma(\frac{n}{2})}{2\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n} \frac{(x_i - y_i) P(y, t)}{\{\mathcal{P}_n(x, y)\}^{\frac{n}{2}}} \prod_{j=1}^n dy_j + f_i - g_i, \quad n \geq 2, \quad x \in \mathbb{R}^n, \quad t \geq 0 \quad (3.6)$$

The difficulty in solving the system of equations (2.1) – (2.3) stems from the presence of the nonlinear term g_i . We therefore recast the Navier-Stokes equation (3.6) as:

$$\frac{\partial v_i}{\partial t} = \kappa \Delta v_i - \mathcal{U}_i(v) + \mathcal{F}_i(x, t) \quad x \in \mathbb{R}^n, \quad t \geq 0 \quad (3.7)$$

where

$$\mathcal{U}_i(x, t) = g_i - \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n} \frac{(x_i - y_i) \sum_{k=1}^n \frac{\partial g_k(y, t)}{\partial y_k}}{\{\mathcal{P}_n(x, y)\}^{\frac{n}{2}}} \prod_{j=1}^n dy_j, \quad x \in \mathbb{R}^n, t \geq 0 \quad (3.8)$$

and

$$\mathcal{F}_i(x, t) = f_i - \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n} \frac{(x_i - w_i) \sum_{k=1}^n \frac{\partial f_k(w, t)}{\partial w_k}}{\{\mathcal{P}_n(x, w)\}^{\frac{n}{2}}} \prod_{j=1}^n dw_j, \quad x \in \mathbb{R}^n, t \geq 0 \quad (3.9)$$

The three terms on the right hand side of (3.7), $\kappa \Delta v_i$, $\mathcal{U}_i(x, t)$ and $\mathcal{F}_i(x, t)$ are associated, respectively, with the linear viscous force, the nonlinear inertial force and the externally applied force acting on the fluid.

A posteriori we state the following *lemma*:

If $\mathcal{U}_i(x, t) \equiv 0$, that is

$$g_i^0(x) = \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} \int_{\mathbb{R}^n} \frac{(x_i - y_i) \sum_{k=1}^n \frac{\partial g_k^0}{\partial y_k}}{\{\mathcal{P}_n(x, y)\}^{\frac{n}{2}}} \prod_{j=1}^n dy_j \quad (3.10)$$

then the solution of the Navier-Stokes equation (2.1) is determined as a solution of the non-homogeneous diffusion equation Carslaw and Jaeger (1959); Thambynayagam (2011):

$$\begin{aligned} v_i(x, t) &= \frac{1}{(2\sqrt{\pi\kappa t})^n} \int_{\mathbb{R}^n} v_i^{(0)}(y) e^{-\sum_{k=1}^n \frac{(x_k - y_k)^2}{4\kappa t}} \prod_{j=1}^n dy_j + \\ &+ \frac{1}{(2\sqrt{\pi\kappa})^n} \int_{\mathbb{R}^n} \int_0^t \frac{\mathcal{F}_i(y, \tau) e^{-\sum_{k=1}^n \frac{(x_k - y_k)^2}{4\kappa(t-\tau)}}}{(t-\tau)^{\frac{n}{2}}} d\tau \prod_{j=1}^n dy_j \end{aligned} \quad (3.11)$$

If the externally applied force on the fluid f_i is set to zero, then, the second term on the right hand side of equation (3.11) vanishes and the solution of the Navier-Stokes equation (2.1) reduces to that of the Cauchy diffusion equation:

$$v_i(x, t) = \frac{1}{(2\sqrt{\pi\kappa t})^n} \int_{\mathbb{R}^n} v_i^{(0)}(y) e^{-\sum_{k=1}^n \frac{(x_k - y_k)^2}{4\kappa t}} \prod_{j=1}^n dy_j \quad (3.12)$$

There are two known solutions given in Appendix B; the Taylor (1923) and the unsteady form of Arnold (1965), which satisfy the relationship (3.10). The integral identities used in this paper are given in Appendix C.

4. A solution in \mathbb{R}^3

We consider $v_i^{(0)}(x)$ of the form

$$\begin{aligned} v_1^0 &= \sin(\alpha x_1 + \xi_1) \cos(\alpha x_2 + \xi_2) \sin(\alpha x_3 + \xi_3) - \\ &\quad - \sin(\alpha x_1 + \xi_2) \cos(\alpha x_3 + \xi_1) \sin(\alpha x_2 + \xi_3) \end{aligned} \quad (4.1)$$

$$\begin{aligned} v_2^0 &= \sin(\alpha x_2 + \xi_1) \cos(\alpha x_3 + \xi_2) \sin(\alpha x_1 + \xi_3) - \\ &\quad - \sin(\alpha x_2 + \xi_2) \cos(\alpha x_1 + \xi_1) \sin(\alpha x_3 + \xi_3) \end{aligned} \quad (4.2)$$

$$\begin{aligned} v_3^0 &= \sin(\alpha x_3 + \xi_1) \cos(\alpha x_1 + \xi_2) \sin(\alpha x_2 + \xi_3) - \\ &\quad - \sin(\alpha x_3 + \xi_2) \cos(\alpha x_2 + \xi_1) \sin(\alpha x_1 + \xi_3) \end{aligned} \quad (4.3)$$

where α is a real number. In (3.10), $g_i^0(x)$ derived from (2.2) can be divided into two parts:

$$g_i^0 = \mathcal{V}_i^0 + \mathcal{W}_i^0 \quad (4.4)$$

\mathcal{V}_i^0 is the sum of all terms that are not functions of x_i . \mathcal{W}_i is the sum of the remaining terms. If we examine closely (3.10), the prerequisite for this relationship to hold is that $\mathcal{V}_i^0 = 0$ exists. The following expressions are obtained:

$$g_1^0(x_1, x_2, x_3) = \mathcal{V}_1^0(x_2, x_3) + \mathcal{W}_1^0(x_1, x_2, x_3) \quad (4.5)$$

$$g_2^0(x_2, x_3, x_1) = \mathcal{V}_2^0(x_3, x_1) + \mathcal{W}_2(x_2, x_3, x_1) \quad (4.6)$$

$$g_3^0(x_3, x_1, x_2) = \mathcal{V}_3^0(x_1, x_2) + \mathcal{W}_3(x_3, x_1, x_2) \quad (4.7)$$

$$\begin{aligned} \mathcal{V}_1^0(x_2, x_3) &= -\frac{\alpha}{4} \cos(\xi_1 - \xi_3) \sin(\xi_2 - \xi_3) \cos(2\alpha x_2 + \xi_2 + \xi_1) - \\ &\quad - \frac{\alpha}{4} \cos(\xi_1 - \xi_3) \cos(\xi_1 - \xi_2) \sin(2\alpha x_3 + \xi_2 + \xi_3) - \\ &\quad - \frac{\alpha}{4} \cos(\xi_3 - \xi_2) \cos(\xi_1 - \xi_2) \sin(2\alpha x_2 + \xi_3 + \xi_1) - \\ &\quad - \frac{\alpha}{4} \cos(\xi_3 - \xi_2) \sin(\xi_1 - \xi_3) \cos(2\alpha x_3 + \xi_1 + \xi_2) + \\ &\quad + \frac{\alpha}{4} \sin(\xi_2 - \xi_1) \sin(\xi_3 - \xi_1) \sin(2\alpha x_2 + \xi_2 + \xi_3) + \\ &\quad + \frac{\alpha}{4} \sin(\xi_2 - \xi_1) \sin(\xi_2 - \xi_3) \sin(2\alpha x_3 + \xi_3 + \xi_1) \end{aligned} \quad (4.8)$$

$$\begin{aligned}
\mathcal{V}_2^0(x_3, x_1) = & -\frac{\alpha}{4} \cos(\xi_1 - \xi_3) \sin(\xi_2 - \xi_3) \cos(2\alpha x_3 + \xi_2 + \xi_1) - \\
& -\frac{\alpha}{4} \cos(\xi_1 - \xi_3) \cos(\xi_1 - \xi_2) \sin(2\alpha x_1 + \xi_2 + \xi_3) - \\
& -\frac{\alpha}{4} \cos(\xi_3 - \xi_2) \cos(\xi_1 - \xi_2) \sin(2\alpha x_3 + \xi_3 + \xi_1) - \\
& -\frac{\alpha}{4} \cos(\xi_3 - \xi_2) \sin(\xi_1 - \xi_3) \cos(2\alpha x_1 + \xi_1 + \xi_2) + \\
& +\frac{\alpha}{4} \sin(\xi_2 - \xi_1) \sin(\xi_3 - \xi_1) \sin(2\alpha x_3 + \xi_2 + \xi_3) + \\
& +\frac{\alpha}{4} \sin(\xi_2 - \xi_1) \sin(\xi_2 - \xi_3) \sin(2\alpha x_1 + \xi_3 + \xi_1) \quad (4.9)
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_3^0(x_1, x_2) = & -\frac{\alpha}{4} \cos(\xi_1 - \xi_3) \sin(\xi_2 - \xi_3) \cos(2\alpha x_1 + \xi_2 + \xi_1) - \\
& -\frac{\alpha}{4} \cos(\xi_1 - \xi_3) \cos(\xi_1 - \xi_2) \sin(2\alpha x_2 + \xi_2 + \xi_3) - \\
& -\frac{\alpha}{4} \cos(\xi_3 - \xi_2) \cos(\xi_1 - \xi_2) \sin(2\alpha x_1 + \xi_3 + \xi_1) - \\
& -\frac{\alpha}{4} \cos(\xi_3 - \xi_2) \sin(\xi_1 - \xi_3) \cos(2\alpha x_2 + \xi_1 + \xi_2) + \\
& +\frac{\alpha}{4} \sin(\xi_2 - \xi_1) \sin(\xi_3 - \xi_1) \sin(2\alpha x_1 + \xi_2 + \xi_3) + \\
& +\frac{\alpha}{4} \sin(\xi_2 - \xi_1) \sin(\xi_2 - \xi_3) \sin(2\alpha x_2 + \xi_3 + \xi_1) \quad (4.10)
\end{aligned}$$

$$\begin{aligned}
\mathcal{W}_1^0(x_1, x_2, x_3) = & \frac{\alpha}{4} \sin(2\alpha x_1 + 2\xi_1) - \frac{\alpha}{4} \sin(2\alpha x_1 + 2\xi_1) \cos(2\alpha x_3 + 2\xi_3) + \\
& +\frac{\alpha}{4} \sin(2\alpha x_1 + 2\xi_2) - \frac{\alpha}{4} \sin(2\alpha x_1 + 2\xi_2) \cos(2\alpha x_2 + 2\xi_3) - \\
& -\frac{\alpha}{2} \sin(\xi_3 - \xi_2) \sin(\xi_3 - \xi_1) \sin(2\alpha x_1 + \xi_1 + \xi_2) - \\
& -\frac{\alpha}{4} \sin(\xi_3 - \xi_1) \sin(2\alpha x_1 + \xi_1 + \xi_2) \sin(2\alpha x_2 + \xi_2 + \xi_3) - \\
& -\frac{\alpha}{4} \sin(\xi_3 - \xi_2) \sin(2\alpha x_1 + \xi_1 + \xi_2) \sin(2\alpha x_3 + \xi_3 + \xi_1) + \\
& +\frac{\alpha}{4} \sin(\xi_2 - \xi_3) \cos(2\alpha x_1 + \xi_3 + \xi_1) \cos(2\alpha x_2 + \xi_1 + \xi_2) + \\
& +\frac{\alpha}{4} \cos(\xi_1 - \xi_2) \cos(2\alpha x_1 + \xi_3 + \xi_1) \sin(2\alpha x_3 + \xi_2 + \xi_3) - \\
& +\frac{\alpha}{4} \cos(\xi_1 - \xi_2) \cos(2\alpha x_1 + \xi_2 + \xi_3) \sin(2\alpha x_2 + \xi_3 + \xi_1) + \\
& +\frac{\alpha}{4} \sin(\xi_1 - \xi_3) \cos(2\alpha x_1 + \xi_2 + \xi_3) \cos(2\alpha x_3 + \xi_1 + \xi_2) \quad (4.11)
\end{aligned}$$

$$\begin{aligned}
 \mathcal{W}_2^0(x_2, x_3, x_1) &= \frac{\alpha}{4} \sin(2\alpha x_2 + 2\xi_1) - \frac{\alpha}{4} \sin(2\alpha x_2 + 2\xi_1) \cos(2\alpha x_1 + 2\xi_3) + \\
 &+ \frac{\alpha}{4} \sin(2\alpha x_2 + 2\xi_2) - \frac{\alpha}{4} \sin(2\alpha x_2 + 2\xi_2) \cos(2\alpha x_3 + 2\xi_3) - \\
 &- \frac{\alpha}{2} \sin(\xi_3 - \xi_2) \sin(\xi_3 - \xi_1) \sin(2\alpha x_2 + \xi_1 + \xi_2) - \\
 &- \frac{\alpha}{4} \sin(\xi_3 - \xi_1) \sin(2\alpha x_2 + \xi_1 + \xi_2) \sin(2\alpha x_3 + \xi_2 + \xi_3) - \\
 &- \frac{\alpha}{4} \sin(\xi_3 - \xi_2) \sin(2\alpha x_2 + \xi_1 + \xi_2) \sin(2\alpha x_1 + \xi_3 + \xi_1) + \\
 &+ \frac{\alpha}{4} \sin(\xi_2 - \xi_3) \cos(2\alpha x_2 + \xi_3 + \xi_1) \cos(2\alpha x_3 + \xi_1 + \xi_2) + \\
 &+ \frac{\alpha}{4} \cos(\xi_1 - \xi_2) \cos(2\alpha x_2 + \xi_3 + \xi_1) \sin(2\alpha x_1 + \xi_2 + \xi_3) - \\
 &+ \frac{\alpha}{4} \cos(\xi_1 - \xi_2) \cos(2\alpha x_2 + \xi_2 + \xi_3) \sin(2\alpha x_3 + \xi_3 + \xi_1) + \\
 &+ \frac{\alpha}{4} \sin(\xi_1 - \xi_3) \cos(2\alpha x_2 + \xi_2 + \xi_3) \cos(2\alpha x_1 + \xi_1 + \xi_2) \quad (4.12)
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{W}_3^0(x_3, x_1, x_2) &= \frac{\alpha}{4} \sin(2\alpha x_3 + 2\xi_1) - \frac{\alpha}{4} \sin(2\alpha x_3 + 2\xi_1) \cos(2\alpha x_2 + 2\xi_3) + \\
 &+ \frac{\alpha}{4} \sin(2\alpha x_3 + 2\xi_2) - \frac{\alpha}{4} \sin(2\alpha x_3 + 2\xi_2) \cos(2\alpha x_1 + 2\xi_3) - \\
 &- \frac{\alpha}{2} \sin(\xi_3 - \xi_2) \sin(\xi_3 - \xi_1) \sin(2\alpha x_3 + \xi_1 + \xi_2) - \\
 &- \frac{\alpha}{4} \sin(\xi_3 - \xi_1) \sin(2\alpha x_3 + \xi_1 + \xi_2) \sin(2\alpha x_1 + \xi_2 + \xi_3) - \\
 &- \frac{\alpha}{4} \sin(\xi_3 - \xi_2) \sin(2\alpha x_3 + \xi_1 + \xi_2) \sin(2\alpha x_2 + \xi_3 + \xi_1) + \\
 &+ \frac{\alpha}{4} \sin(\xi_2 - \xi_3) \cos(2\alpha x_3 + \xi_3 + \xi_1) \cos(2\alpha x_1 + \xi_1 + \xi_2) + \\
 &+ \frac{\alpha}{4} \cos(\xi_1 - \xi_2) \cos(2\alpha x_3 + \xi_3 + \xi_1) \sin(2\alpha x_2 + \xi_2 + \xi_3) - \\
 &+ \frac{\alpha}{4} \cos(\xi_1 - \xi_2) \cos(2\alpha x_3 + \xi_2 + \xi_3) \sin(2\alpha x_1 + \xi_3 + \xi_1) + \\
 &+ \frac{\alpha}{4} \sin(\xi_1 - \xi_3) \cos(2\alpha x_3 + \xi_2 + \xi_3) \cos(2\alpha x_2 + \xi_1 + \xi_2) \quad (4.13)
 \end{aligned}$$

We note that $\xi_1 = -\frac{\pi}{3}$, $\xi_2 = \frac{\pi}{3}$, and $\xi_3 = \frac{\pi}{2}$ renders $\mathcal{V}_i^0 = 0$, and g_i^0 equals \mathcal{W}_i^0 .

The resulting expressions for g_i^0 and $\sum_{i=1}^3 \frac{\partial g_i^0}{\partial x_i}$ are:

$$\begin{aligned}
 g_1^0(x_1, x_2, x_3) &= -\frac{3\alpha}{8} \sin(2\alpha x_1) - \frac{3\alpha}{32} \sin(2\alpha x_1) \cos(2\alpha x_2) - \frac{3\alpha}{32} \sin(2\alpha x_1) \cos(2\alpha x_3) - \\
 &- \frac{3\sqrt{3}\alpha}{32} \cos(2\alpha x_1) \cos(2\alpha x_3) + \frac{3\sqrt{3}\alpha}{32} \cos(2\alpha x_1) \cos(2\alpha x_2) + \\
 &+ \frac{3\sqrt{3}\alpha}{32} \sin(2\alpha x_1) \sin(2\alpha x_2) - \frac{3\sqrt{3}\alpha}{32} \sin(2\alpha x_1) \sin(2\alpha x_3) + \\
 &+ \frac{3\alpha}{32} \cos(2\alpha x_1) \sin(2\alpha x_3) + \frac{3\alpha}{32} \cos(2\alpha x_1) \sin(2\alpha x_2) \quad (4.14)
 \end{aligned}$$

$$\begin{aligned}
g_2^0(x_2, x_3, x_1) &= -\frac{3\alpha}{8} \sin(2\alpha x_2) - \frac{3\alpha}{32} \sin(2\alpha x_2) \cos(2\alpha x_3) - \frac{3\alpha}{32} \sin(2\alpha x_2) \cos(2\alpha x_1) - \\
&\quad - \frac{3\sqrt{3}\alpha}{32} \cos(2\alpha x_2) \cos(2\alpha x_1) + \frac{3\sqrt{3}\alpha}{32} \cos(2\alpha x_2) \cos(2\alpha x_3) + \\
&\quad + \frac{3\sqrt{3}\alpha}{32} \sin(2\alpha x_2) \sin(2\alpha x_3) - \frac{3\sqrt{3}\alpha}{32} \sin(2\alpha x_2) \sin(2\alpha x_1) + \\
&\quad + \frac{3\alpha}{32} \cos(2\alpha x_2) \sin(2\alpha x_1) + \frac{3\alpha}{32} \cos(2\alpha x_2) \sin(2\alpha x_3) \tag{4.15}
\end{aligned}$$

$$\begin{aligned}
g_3^0(x_3, x_1, x_2) &= -\frac{3\alpha}{8} \sin(2\alpha x_3) - \frac{3\alpha}{32} \sin(2\alpha x_3) \cos(2\alpha x_1) - \frac{3\alpha}{32} \sin(2\alpha x_3) \cos(2\alpha x_2) - \\
&\quad - \frac{3\sqrt{3}\alpha}{32} \cos(2\alpha x_3) \cos(2\alpha x_2) + \frac{3\sqrt{3}\alpha}{32} \cos(2\alpha x_3) \cos(2\alpha x_1) + \\
&\quad + \frac{3\sqrt{3}\alpha}{32} \sin(2\alpha x_3) \sin(2\alpha x_1) - \frac{3\sqrt{3}\alpha}{32} \sin(2\alpha x_3) \sin(2\alpha x_2) + \\
&\quad + \frac{3\alpha}{32} \cos(2\alpha x_3) \sin(2\alpha x_2) + \frac{3\alpha}{32} \cos(2\alpha x_3) \sin(2\alpha x_1) \tag{4.16}
\end{aligned}$$

$$\begin{aligned}
\sum_{i=1}^3 \frac{\partial g_i^0}{\partial x_i} &= -\frac{3\alpha^2}{4} \cos(2\alpha x_1) - \frac{3\alpha^2}{4} \cos(2\alpha x_2) - \frac{3\alpha^2}{4} \cos(2\alpha x_3) - \\
&\quad - \frac{3\alpha^2}{8} \cos(2\alpha x_1) \cos(2\alpha x_2) - \frac{3\alpha^2}{8} \cos(2\alpha x_1) \cos(2\alpha x_3) + \\
&\quad + \frac{3\sqrt{3}\alpha^2}{8} \sin(2\alpha x_1) \cos(2\alpha x_3) - \frac{3\sqrt{3}\alpha^2}{8} \sin(2\alpha x_1) \cos(2\alpha x_2) + \\
&\quad + \frac{3\sqrt{3}\alpha^2}{8} \cos(2\alpha x_1) \sin(2\alpha x_2) - \frac{3\sqrt{3}\alpha^2}{8} \cos(2\alpha x_1) \sin(2\alpha x_3) - \\
&\quad - \frac{3\alpha^2}{8} \sin(2\alpha x_1) \sin(2\alpha x_3) - \frac{3\alpha^2}{8} \sin(2\alpha x_1) \sin(2\alpha x_2) - \\
&\quad - \frac{3\alpha^2}{8} \cos(2\alpha x_2) \cos(2\alpha x_3) - \frac{3\sqrt{3}\alpha^2}{8} \sin(2\alpha x_2) \cos(2\alpha x_3) + \\
&\quad + \frac{3\sqrt{3}\alpha^2}{8} \cos(2\alpha x_2) \sin(2\alpha x_3) - \frac{3\alpha^2}{8} \sin(2\alpha x_2) \sin(2\alpha x_3) \tag{4.17}
\end{aligned}$$

In (3.10), substituting for $\sum_{i=1}^3 \frac{\partial g_i^0}{\partial x_i}$, and performing term by term the integrals on the right hand side, we show that the solution satisfies the necessary condition $\mathcal{U}_i(x, t) \equiv 0$.

Substituting for the initial condition $v_i^{(0)}(x)$ with $\xi_1 = -\frac{\pi}{3}$, $\xi_2 = \frac{\pi}{3}$, and $\xi_3 = \frac{\pi}{2}$ in (3.12) and performing the integrations, we arrive at the solution of the Navier-Stokes equation (2.1):

$$v_1 = \left[\sin \left(\alpha x_1 - \frac{\pi}{3} \right) \cos \left(\alpha x_2 + \frac{\pi}{3} \right) \sin \left(\alpha x_3 + \frac{\pi}{2} \right) - \sin \left(\alpha x_1 + \frac{\pi}{3} \right) \cos \left(\alpha x_3 - \frac{\pi}{3} \right) \sin \left(\alpha x_2 + \frac{\pi}{2} \right) \right] e^{-3\alpha^2 \kappa t} \quad (4.18)$$

$$v_2 = \left[\sin \left(\alpha x_2 - \frac{\pi}{3} \right) \cos \left(\alpha x_3 + \frac{\pi}{3} \right) \sin \left(\alpha x_1 + \frac{\pi}{2} \right) - \sin \left(\alpha x_2 + \frac{\pi}{3} \right) \cos \left(\alpha x_1 - \frac{\pi}{3} \right) \sin \left(\alpha x_3 + \frac{\pi}{2} \right) \right] e^{-3\alpha^2 \kappa t} \quad (4.19)$$

$$v_3 = \left[\sin \left(\alpha x_3 - \frac{\pi}{3} \right) \cos \left(\alpha x_1 + \frac{\pi}{3} \right) \sin \left(\alpha x_2 + \frac{\pi}{2} \right) - \sin \left(\alpha x_3 + \frac{\pi}{3} \right) \cos \left(\alpha x_2 - \frac{\pi}{3} \right) \sin \left(\alpha x_1 + \frac{\pi}{2} \right) \right] e^{-3\alpha^2 \kappa t} \quad (4.20)$$

We may also express (4.18)–(4.20) as:

$$v_1 = \frac{3}{4} \left[\cos(\alpha x_1) \sin(\alpha x_2 - \alpha x_3) - \frac{1}{\sqrt{3}} \sin(\alpha x_1) \sin(\alpha x_2 + \alpha x_3) - \frac{2}{\sqrt{3}} \cos(\alpha x_1) \cos(\alpha x_2) \cos(\alpha x_3) \right] e^{-3\alpha^2 \kappa t} \quad (4.21)$$

$$v_2 = \frac{3}{4} \left[\cos(\alpha x_2) \sin(\alpha x_3 - \alpha x_1) - \frac{1}{\sqrt{3}} \sin(\alpha x_2) \sin(\alpha x_3 + \alpha x_1) - \frac{2}{\sqrt{3}} \cos(\alpha x_2) \cos(\alpha x_3) \cos(\alpha x_1) \right] e^{-3\alpha^2 \kappa t} \quad (4.22)$$

$$v_3 = \frac{3}{4} \left[\cos(\alpha x_3) \sin(\alpha x_1 - \alpha x_2) - \frac{1}{\sqrt{3}} \sin(\alpha x_3) \sin(\alpha x_1 + \alpha x_2) - \frac{2}{\sqrt{3}} \cos(\alpha x_3) \cos(\alpha x_1) \cos(\alpha x_2) \right] e^{-3\alpha^2 \kappa t} \quad (4.23)$$

Pressure is obtained from the solution of the Poisson equation (3.2). Because (3.10) is satisfied, we can express (3.5) as follows:

$$\frac{\partial p}{\partial x_i} = \rho g_i \quad i = 1, 2, 3 \quad (4.24)$$

This leads to straightforward integration for pressure:

$$\begin{aligned} p = & \frac{3\rho}{16} [\cos(2\alpha x_1) + \cos(2\alpha x_2) + \cos(2\alpha x_3)] e^{-6\alpha^2 \kappa t} + \\ & + \frac{3\rho}{64} [\cos(2\alpha x_1 - 2\alpha x_2) + \cos(2\alpha x_3 - 2\alpha x_1)] e^{-6\alpha^2 \kappa t} + \\ & + \frac{3\sqrt{3}\rho}{64} [\sin(2\alpha x_1 - 2\alpha x_2) + \sin(2\alpha x_3 - 2\alpha x_1) + \\ & + \sin(2\alpha x_2 - 2\alpha x_3) + \cos(2\alpha x_3 - 2\alpha x_2)] e^{-6\alpha^2 \kappa t} \end{aligned} \quad (4.25)$$

5. Concluding Remarks

At the outset, the solution we have presented for the velocity vector field appears to be the same as that of Antuono (2020): (A1)-(A3) except for the augmenting coefficient $\left(\frac{4\sqrt{2}}{3\sqrt{3}}\right)v_0$. However, they are not the same.

Antuono (2020) describes the velocity vector field in a tours $\mathbb{T}^3 = [0, L]^3$, in terms of a wave number divided into two parts, characterized by positive and negative helicity respectively. In his work, a singularity that arose from the derivation was eliminated by imposing the bounded energy constraint (2.5). The velocity components were normalized by a reference velocity v_0 so that the average kinetic energy per unit of mass is $\left(\frac{v_0^2}{2}\right)$ at $t = 0$. The pressure field was obtained in terms of a reference pressure and reference density from the velocity field.

In three dimensions, we have four unknowns, three components of the velocity vector field and pressure, and four equations. It is important to note that defining pressure at the initial time independent of velocity would render the problem ill-posed; therefore, only a solenoidal initial velocity vector field is prescribed. In a Cartesian frame of reference, we have solved the fundamental equations of momentum, the Navier-Stokes equation and mass, the continuity equation for incompressible fluids. The method we have presented does not necessitate the imposition of the bounded energy constraint (2.5). The solutions (4.18)-(4.20) and (4.21) for velocity and pressure, respectively, are valid in the whole unbounded domain $\mathbb{R}^3 = \{-\infty < x_i < \infty; i = 1, 2, 3\}$. We have shown that the solutions of the Navier-Stokes system of equations for incompressible fluids for the velocity field and pressure are given by the Cauchy diffusion equation and the Poisson equation, respectively, provided the condition (3.10) is satisfied.

Appendix A. Solution with positive helicity given by Antuono (2020)

$$v_1 = \left(\frac{4\sqrt{2}}{3\sqrt{3}}\right)v_0 \left[\sin\left(\omega x_1 - \frac{\pi}{3}\right) \cos\left(\omega x_2 + \frac{\pi}{3}\right) \sin\left(\omega x_3 + \frac{\pi}{2}\right) - \sin\left(\omega x_1 + \frac{\pi}{3}\right) \cos\left(\omega x_3 - \frac{\pi}{3}\right) \sin\left(\omega x_2 + \frac{\pi}{2}\right) \right] e^{-3\omega^2 \kappa t} \quad (\text{A } 1)$$

$$v_2 = \left(\frac{4\sqrt{2}}{3\sqrt{3}}\right)v_0 \left[\sin\left(\omega x_2 - \frac{\pi}{3}\right) \cos\left(\omega x_3 + \frac{\pi}{3}\right) \sin\left(\omega x_1 + \frac{\pi}{2}\right) - \sin\left(\omega x_2 + \frac{\pi}{3}\right) \cos\left(\omega x_1 - \frac{\pi}{3}\right) \sin\left(\omega x_3 + \frac{\pi}{2}\right) \right] e^{-3\omega^2 \kappa t} \quad (\text{A } 2)$$

$$v_3 = \left(\frac{4\sqrt{2}}{3\sqrt{3}} \right) v_0 \left[\sin \left(\omega x_3 - \frac{\pi}{3} \right) \cos \left(\omega x_1 + \frac{\pi}{3} \right) \sin \left(\omega x_2 + \frac{\pi}{2} \right) - \sin \left(\omega x_3 + \frac{\pi}{3} \right) \cos \left(\omega x_2 - \frac{\pi}{3} \right) \sin \left(\omega x_1 + \frac{\pi}{2} \right) \right] e^{-3\omega^2 \kappa t} \quad (\text{A } 3)$$

where v_1, v_2 and v_3 are the components of the velocity vector field, v_0 is the reference velocity, κ is the kinematical viscosity and ω is the wave number.

Appendix B. Derivations of two known solutions Taylor (1923); Arnold (1965) satisfying equation (3.10) presented from Thambynayagam (2013)

B.1. *The Taylor solution in $\mathbb{R}^2 = \{-\infty < x_i < \infty; i = 1, 2\}$*

$$v_1^{(0)} = \sin(\pi x_1) \cos(\pi x_2), v_2^{(0)} = -\cos(\pi x_1) \sin(\pi x_2) \text{ and } f_1 = f_2 = 0.$$

Substituting for $v_1^{(0)}$ and $v_2^{(0)}$ in (2.3) and (3.10) we obtain

$$g_i^{(0)} = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{(x_i - y_i) \sum_{k=1}^2 \frac{\partial g_k^{(0)}(y,t)}{\partial y_k}}{\mathcal{P}_2(x,y)} \prod_{j=1}^n dy_j = \frac{\pi}{2} \sin(2\pi x_i) \quad (\text{B } 1)$$

resulting in $\mathcal{U}_i(x,t) \equiv 0$. $v(x,t), p(x,t)$ are obtained from (3.11) and (3.2):

$$v_1 = \sin(\pi x_1) \cos(\pi x_2) e^{-2\pi^2 \kappa t} \quad (\text{B } 2)$$

$$v_2 = -\cos(\pi x_1) \sin(\pi x_2) e^{-2\pi^2 \kappa t} \quad (\text{B } 3)$$

and

$$p = -\frac{\rho e^{-4\pi^2 \kappa t}}{4} [\cos(2\pi x_1) + \cos(2\pi x_2)] \quad (\text{B } 4)$$

B.2. *Arnold-Beltrami-Childress (ABC) flows in $\mathbb{R}^3 = \{-\infty < x_i < \infty; i = 1, 2, 3\}$*

$$v_1^{(0)} = a \sin \pi x_3 - c \cos \pi x_2, v_2^{(0)} = b \sin \pi x_1 - a \cos \pi x_3, v_3^{(0)} = c \sin \pi x_2 - b \cos \pi x_1 \text{ and } f_1 = f_2 = f_3 = 0. a, b \text{ and } c \text{ are real constants.}$$

Substituting for $v_1^{(0)}, v_2^{(0)}$ and $v_3^{(0)}$ in (2.3) and (3.10) we obtain

$$g_1^{(0)} = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{(x_1 - y_1) \sum_{k=1}^3 \frac{\partial g_k^{(0)}(y,t)}{\partial y_k}}{\{\mathcal{P}_3(x,y)\}^{\frac{3}{2}}} \prod_{j=1}^3 dy_j = \pi \{bc \sin(\pi x_1) \sin(\pi x_2) - ab \cos(\pi x_1) \cos(\pi x_3)\} \quad (\text{B } 5)$$

$$\begin{aligned}
g_2^{(0)} &= \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{(x_2 - y_2) \sum_{k=1}^3 \frac{\partial g_k^{(0)}(y,t)}{\partial y_k}}{\{\mathcal{P}_3(x,y)\}^{\frac{3}{2}}} \prod_{j=1}^3 dy_j \\
&= \pi \{ac \sin(\pi x_2) \sin(\pi x_3) - bc \cos(\pi x_1) \cos(\pi x_2)\}
\end{aligned} \tag{B 6}$$

$$\begin{aligned}
g_3^{(0)} &= \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{(x_3 - y_3) \sum_{k=1}^3 \frac{\partial g_k^{(0)}(y,t)}{\partial y_k}}{\{\mathcal{P}_3(x,y)\}^{\frac{3}{2}}} \prod_{j=1}^3 dy_j \\
&= \pi \{ab \sin(\pi x_1) \sin(\pi x_3) - ac \cos(\pi x_2) \cos(\pi x_3)\}
\end{aligned} \tag{B 7}$$

$v(x, t)$, $p(x, t)$ are obtained from (3.11) and (3.2):

$$v_1 = \{a \sin(\pi x_3) - c \cos(\pi x_2)\} e^{-\pi^2 \kappa t} \tag{B 8}$$

$$v_2 = \{b \sin(\pi x_1) - a \cos(\pi x_3)\} e^{-\pi^2 \kappa t} \tag{B 9}$$

$$v_3 = \{c \sin(\pi x_2) - b \cos(\pi x_1)\} e^{-\pi^2 \kappa t} \tag{B 10}$$

and

$$p = -\rho e^{-2\pi^2 \kappa t} [bc \cos(\pi x_1) \sin(\pi x_2) + ab \cos(\pi x_3) \sin(\pi x_1) + ac \cos(\pi x_2) \sin(\pi x_3)] \tag{B 11}$$

Appendix C. Integral identities used in this paper

$$\int_{-\infty}^{\infty} e^{-\frac{(x-u)^2}{4\tau}} du = 2\sqrt{\pi\tau} \tag{C 1}$$

$$\int_{-\infty}^{\infty} \sin(\alpha u) e^{-\frac{(x-u)^2}{4\tau}} du = 2\sqrt{\pi\tau} e^{-\alpha^2 \tau} \sin(\alpha x) \tag{C 2}$$

$$\int_{-\infty}^{\infty} \cos(\alpha u) e^{-\frac{(x-u)^2}{4\tau}} du = 2\sqrt{\pi\tau} e^{-\alpha^2 \tau} \cos(\alpha x) \tag{C 3}$$

$$\int_0^{\infty} \int_0^{\infty} \int_0^{\infty} \frac{u_1 \cos\{\beta(x_1 - u_1)\}}{[u_1^2 + u_2^2 + u_3^2]^{\frac{3}{2}}} du_1 du_2 du_3 = \frac{4\pi}{\beta} \sin(\beta x_1) \tag{C 4}$$

$$\int_0^{\infty} \int_0^{\infty} \int_0^{\infty} \frac{u_1 \cos\{\beta(x_1 - u_1)\} \cos\{\beta(x_2 - u_2)\}}{[u_1^2 + u_2^2 + u_3^2]^{\frac{3}{2}}} du_1 du_2 du_3 = \frac{2\pi}{\beta} \sin(\beta x_1) \cos(\beta x_2) \tag{C 5}$$

$$\int_0^\infty \int_0^\infty \int_0^\infty \frac{u_1 \sin\{\beta(x_1 - u_1)\} \cos\{\beta(x_2 - u_2)\}}{[u_1^2 + u_2^2 + u_3^2]^{\frac{3}{2}}} du_1 du_2 du_3 = -\frac{2\pi}{\beta} \cos(\beta x_1) \cos(\beta x_2) \quad (\text{C } 6)$$

$$\int_0^\infty \int_0^\infty \int_0^\infty \frac{u_1 \cos\{\beta(x_1 - u_1)\} \sin\{\beta(x_2 - u_2)\}}{[u_1^2 + u_2^2 + u_3^2]^{\frac{3}{2}}} du_1 du_2 du_3 = \frac{2\pi}{\beta} \sin(\beta x_1) \sin(\beta x_2) \quad (\text{C } 7)$$

$$\int_0^\infty \int_0^\infty \int_0^\infty \frac{u_1 \sin\{\beta(x_1 - u_1)\} \sin\{\beta(x_2 - u_2)\}}{[u_1^2 + u_2^2 + u_3^2]^{\frac{3}{2}}} du_1 du_2 du_3 = -\frac{2\pi}{\beta} \cos(\beta x_1) \sin(\beta x_2) \quad (\text{C } 8)$$

REFERENCES

- Antuono, M. (2020). *J. Fluid. Mech* 890, A23.
- Arnold, V. I. (1965). *C. R. Acad. Sci. Paris* 261, 17–20.
- Batchelor, G. K. (1967). *An Introduction to Fluid Dynamics*. Cambridge: Cambridge University Press.
- Carslaw, H. S. and J. C. Jaeger (1959). *Conduction of Heat in Solids*. Oxford: Clarendon Press.
- Drazin, P. and N. Riley (2007). *The Navier-Stokes Equations: a classification of flows and exact solutions*. Cambridge: Cambridge University Press.
- Ethier, C. R. and D. A. Steinman (1994). *Int. J. Numer. Mecth. Fluids* 19, 369–375.
- Gresho, P. M. and R. L. Sani (1987). On pressure boundary conditions for the incompressible navier-stokes equations. *International Journal for Numerical Methods in Fluids*, 7, 1111–1145.
- Ladyzhenskaya, O. (1969). *The Mathematical Theory of Viscous Incompressible Flows*. New York: Gordon and Breach.
- Navier, C. L. (1827). Memoire sur less lois du mouvement des fluides. *Mem. Acad. Sci. Inst. France* 6, 389–440.
- Stokes, G. G. (1845). On the theories of the internal friction of fluids in motion, and of the equilibrium and motion of elastic solids. *Trans. Cambridge Philos. Soc.* 8, 287–305.
- Taylor, G. I. (1923). On the decay of vortices in a viscous fluid. *Philos. Mag.* 46, 671–674.
- Thambynayagam, R. K. M. (2011). *The Diffusion Handbook: Applied Solutions for Engineers*. New York: McGraw-Hill.
- Thambynayagam, R. K. M. (2013). Classical analytic solutions of the non-stationary navier-stokes equation in two, three and higher dimensions. *arXiv:1307.7632 [math-ph]*.